

# Protein kinase C activity blocks neuropeptide Y-mediated inhibition of glutamate release and contributes to excitability of the hippocampus in status epilepticus

Ana P. Silva,<sup>\*,‡</sup> Joana Lourenço,<sup>‡</sup> Sara Xapelli,<sup>‡</sup> Raquel Ferreira,<sup>‡</sup> Heidi Kristiansen,<sup>§</sup> David P. D. Woldbye,<sup>§</sup> Catarina R. Oliveira,<sup>†,‡</sup> and João O. Malva<sup>†,‡,1</sup>

<sup>\*</sup>Institute of Pharmacology and Therapeutics and <sup>†</sup>Institute of Biochemistry, Faculty of Medicine, University of Coimbra, Coimbra, Portugal; <sup>‡</sup>Center for Neuroscience and Cell Biology, University of Coimbra, Coimbra, Portugal; <sup>§</sup>Laboratory of Neuropsychiatry, Rigshospitalet University Hospital and Department of Pharmacology, University of Copenhagen, Denmark

**ABSTRACT** The unbalanced excitatory/inhibitory neurotransmitter function in the neuronal network afflicted by seizures is the main biochemical and biophysical hallmark of epilepsy. The aim of this work was to identify changes in the signaling mechanisms associated with neuropeptide Y (NPY)-mediated inhibition of glutamate release that may contribute to hyperexcitability. Using isolated rat hippocampal nerve terminals, we showed that the KCl-evoked glutamate release is inhibited by NPY Y<sub>2</sub> receptor activation and is potentiated by the stimulation of protein kinase C (PKC). Moreover, we observed that immediately after status epilepticus (6 h postinjection with kainate, 10 mg/kg), the functional inhibition of glutamate release by NPY Y<sub>2</sub> receptors was transiently blocked concomitantly with PKC hyperactivation. The pharmacological blockade of seizure-activated PKC revealed again the Y<sub>2</sub> receptor-mediated inhibition of glutamate release. The functional activity of PKC immediately after status epilepticus was assessed by evaluating phosphorylation of the AMPA receptor subunit GluR1 (Ser-831), a substrate for PKC. Moreover, NPY-stimulated [<sup>35</sup>S]GTPγS autoradiographic binding studies indicated that the common target for Y<sub>2</sub> receptor and PKC on the inhibition/potential of glutamate release was located downstream of the Y<sub>2</sub> receptor, or its interacting G-protein, and involves voltage-gated calcium channels.—Silva, A. P., Lourenço, J., Xapelli, S., Ferreira, R., Kristiansen, H., Woldbye, D. P. D., Oliveira, C. R., Malva, J. O. Protein kinase C activity blocks neuropeptide Y-mediated inhibition of glutamate release and contributes to excitability of the hippocampus in status epilepticus. *FASEB J.* 21, 671–681 (2007)

**Key Words:** Y<sub>2</sub> receptor • nerve terminal • seizures • phosphorylation

NEUROPEPTIDE Y (NPY) HAS BEEN IMPLICATED in several centrally mediated physiological functions such as reg-

ulation of circadian rhythms, body temperature, sexual behavior, blood pressure, appetite, and neuroendocrine secretions (1–3). This peptide has also been shown to modulate anxiety-related disorders and cognitive functions such as learning and memory (4), and it has been widely suggested to be involved in epileptogenesis and epilepsy (5–7). Moreover, the broad physiological actions of NPY are transduced by at least five cloned NPY receptors subtypes (Y<sub>1</sub>, Y<sub>2</sub>, Y<sub>4</sub>, Y<sub>5</sub>, and y<sub>6</sub>), which belong to the G-protein-coupled receptor superfamily (8). NPY and NPY Y<sub>1</sub>, Y<sub>2</sub>, and Y<sub>5</sub> receptors are particularly abundant in the hippocampus, where they play a major role in regulating glutamate release (3, 9).

Endogenous NPY seems to play a major role in regulating seizure activity, and this is supported by the massive increase of NPY levels under epileptic conditions in both inhibitory interneurons and excitatory granule cells (5, 10). Moreover, seizure-related increase in NPY expression is accompanied by modified levels of NPY receptor subtypes in the hippocampus (11), and it seems that more than one receptor subtype could be responsible for mediating the antiepileptic effects of NPY, but clearly Y<sub>2</sub> and/or Y<sub>5</sub> receptors are key players (10, 12, 13). It was reported that NPY-deficient mice have increased susceptibility to seizures induced by pentylenetetrazol or kainate and that seizure activity was reduced in rat models of NPY overexpression (14–17). Furthermore, NPY acting *via* Y<sub>2</sub> receptors can inhibit excitatory synaptic transmission (18, 19), the release of glutamate (20, 21), and epileptiform activity (10, 22) in hippocampal slices. NPY was also found to inhibit kainate-induced seizures *via* Y<sub>5</sub> receptors (13, 23), but recently El Bahh and collaborators (12) demonstrated that Y<sub>2</sub> receptors play a key role in the

<sup>1</sup>Correspondence: Center for Neuroscience and Cell Biology, Institute of Biochemistry, Faculty of Medicine, University of Coimbra, 3004–504 Coimbra, Portugal. E-mail: jomalva@fmed.uc.pt  
doi: 10.1096/fj.06-6163com

antiepileptic properties of NPY in the hippocampus. Recently, we observed that in different stages of the epileptogenic process, the ability of different NPY receptors to modulate the release of glutamate is changed (21). We suggested that in the acute phase of epilepsy, a transient loss of NPY-mediated inhibition of glutamate release might be associated with hyperexcitability of the hippocampal neuronal network and seizure activity. After the acute phase, an adaptation of the NPYergic modulation of glutamate release was identified and the inhibitory effect of NPY was again close to control levels (21).

It is well established that protein kinase A- and C-mediated substrate phosphorylation is implicated in a broad spectrum of neuronal events, including neuronal differentiation, process outgrowth, axonal regeneration (24–26), glutamate-induced neurotoxicity (27), and neurotransmitter release (28, 29). It has also been shown that protein kinases act on exocytotic machinery (30) and modulate ion channels (29, 31) and receptor activity/function (32).

Taken together, the current data indicate that NPY receptors clearly play a role in normal physiological conditions, as well as in response to pathological hyperactivity in the hippocampus (6, 21, 33, 34). Yet despite the role of these receptors already mentioned, there is little consensus about the intracellular signaling pathway (or pathways) that can interfere with their function under pathological conditions such as epilepsy.

## MATERIALS AND METHODS

### Rodent model of temporal lobe epilepsy

Male Wistar rats 6–7 wk of age were used. Kainate (KA; Ocean Produce International, Shelburne, Nova Scotia, Canada) was dissolved in a maximum volume of 500  $\mu$ l of sterile 0.9% NaCl and injected intraperitoneally (i.p.) at a dosage of 10 mg/kg body wt. The control group of animals was injected with the same volume of saline (21). Rats were observed for at least 3 h and their behavior was rated as described previously (35). All the animals used in the present work achieved status epilepticus and were sacrificed 6 h or 24 h after KA injection.

All procedures involving experimental animals were performed in accordance with European Community guidelines. All efforts were made to minimize animal suffering and to reduce the number of animals used.

### Preparation of rat hippocampal synaptosomes

A partially purified synaptosomal fraction ( $P_2$ ) was isolated from hippocampi or from hippocampal subregions CA1, CA3, and dentate gyrus (DG) of male Wistar control or epileptic rats (sacrificed 6 h or 24 h after KA injection), essentially as described for brain cortex (36), with some modifications (21, 37, 38). The hippocampi were homogenized in 0.32 M sucrose, 10 mM HEPES-Na, pH 7.4, using a Thomas B-Potter homogenizer (Thomas Scientific, Swedesboro, NJ, USA) and centrifuged at 3000 *g* for 2 min. The pellet obtained was resuspended, followed by sedimentation at the same speed. The combined supernatants were spun for 12 min at 14,600 *g* and a  $P_2$  pellet was obtained. The upper

white layer of the pellet containing synaptosomes (39) was removed with a small spoon and resuspended in the same sucrose medium used before.

Coronal slices of hippocampus (800  $\mu$ m-thick) were prepared for isolation of the synaptosomes from hippocampal subregions (CA1, CA3, and DG) of nonepileptic and epileptic rats (6 h and 24 h after KA injection). In each slice, the fimbria and the subiculum were separated from the rest of the slice under stereomicroscopic observation. CA3 subslices were obtained by separation from CA1 and DG, and the last separation (CA1 from DG) was performed through the hippocampal sulcus (38). The pooled subslices were homogenized in the sucrose medium using a Thomas AA-Potter, transferred to Eppendorf tubes, and centrifuged as described for the isolation of whole hippocampal synaptosomes. The protein concentration was determined by the Biuret method (40) for glutamate release experiments, and the synaptosomes were stored as drained pellets containing 1 mg of protein.

### Preparation of hippocampal lysates

For Western blot analysis, the hippocampi or hippocampal subregions (CA1, CA3, and DG) of male Wistar control or epileptic rats (sacrificed 6 h or 24 h after KA injection) were removed as described in the foregoing section. Then the tissue was individually homogenized with 40 strokes in a glass homogenizer in 50 mM Tris-HCl, 0.5% Triton X-100, supplemented with 100  $\mu$ M phenylmethanesulfonyl fluoride, 1 mM dithiothreitol (DTT), 1  $\mu$ g/ml chymostatin, 1  $\mu$ g/ml leupeptin, 1  $\mu$ g/ml antipain, 5  $\mu$ g/ml pepstatin A (all from Sigma Chemical Co., St. Louis, MO, USA), pH 7.4, at 4°C. Protein concentration was determined by the bicinchoninic acid method (41).

### Measurement of glutamate release

The release of endogenous glutamate was followed using a continuous fluorimetric assay as described previously (42), with some modifications (37, 38). Synaptosomes (1 mg protein) were preincubated for 20 min at 37°C in the following medium (in mM): 132 NaCl, 1 KCl, 1 MgCl<sub>2</sub>, 1.2 H<sub>3</sub>PO<sub>4</sub>, 0.1 CaCl<sub>2</sub>, 10 glucose, 10 HEPES-Na, pH 7.4, with 0.1% fatty acid-free BSA. After this period, NPY or Y<sub>1</sub>, Y<sub>2</sub>, or Y<sub>5</sub> receptor agonists ([Leu<sup>31</sup>, Pro<sup>34</sup>]NPY, NPY(13–36) or NPY(19–23)-(Gly<sup>1</sup>, Ser<sup>3</sup>, Gln<sup>4</sup>, Thr<sup>6</sup>, AL<sup>31</sup>, Aib<sup>32</sup>, Gln<sup>34</sup>)-PP, respectively) (Bachem AG, Bubendorf, Switzerland), antagonists [BIBP3226 (Peninsula Labs, Belmont, CA, USA), BIIE0246 (kindly provided by Dr. Henri Doods, Boehringer Ingelheim Pharma KG, Germany) and L-152,804 (Tocris, Bristol, UK), respectively], and/or PKA and PKC activator/inhibitor [8-Br-cAMP/H-89 and phorbol myristate acetate (PMA)/bisindolylmaleimide I (BIS), respectively] (all from Sigma Chemical Co.), were added to the incubation medium for an additional 10 min. Synaptosomes were then centrifuged at 15,800 *g* and resuspended in 1 ml of the same medium without BSA containing 1 mM CaCl<sub>2</sub>. The suspension was transferred to a stirred acrylic cuvette, maintained at 37°C, followed by the addition of 1 mM NADP<sup>+</sup>, 50 U of purified glutamate dehydrogenase, and again NPY receptor agonists or antagonists. Fluorescence was measured using a Perkin-Elmer model LS-5B luminescence spectrometer (Perkin-Elmer Ltd, Beaconsfield, Buckinghamshire, England) at the excitation and emission wavelengths of 340 nm and 460 nm, respectively, with excitation and emission slits of 5 nm and 10 nm, respectively. The data were collected at 0.5 s intervals and at the end of each experiment, 2.5 nmol of L-glutamate was

added as a calibration to allow quantification of released glutamate.

Glutamate release was monitored for 11 min and synaptosomes were stimulated 4 min after the beginning of each experiment with 15 mM KCl or 5  $\mu$ M ionomycin.

### Western blot analysis

Twenty-five micrograms of protein from total hippocampus or from hippocampal subregions were separated by SDS-PAGE on 7.5% acrylamide/bisacrilamide gels, using a Bicine/SDS-based electrophoresis buffer (pH 8.3), and transferred onto PVDF membranes (750 mA, 50 min at 4°C in a solution containing 10 mM CAPS and 10% methanol, pH 11.0) (43). Membrane blocking was performed for 1 h at room temperature in Tris-buffered saline containing 5% low-fat milk and 0.1% Tween 20. Primary antibodies raised against GluR1 and phospho-GluR1 (Ser-831) (both from Upstate, Barcelona, Spain) were applied overnight at 4°C and were detected using alkaline phosphatase conjugated secondary antibodies. Immunoblots were visualized using the Enhanced ChemiFluorescence detection reagent and a Versa Doc 3000 imaging system (Bio-Rad, Hercules, CA, USA).

### Immunohistochemistry

Rats were deeply anesthetized with sodium pentobarbital (120 mg/kg, i.p.) and perfused transcardially first with PBS, then fixed with 4% paraformaldehyde in PBS. Brains were removed and coronal sections (20  $\mu$ m) were cut on a cryostat. Brain sections placed on gelatin-coated glass slides were washed with 0.05 M Tris-buffered saline (TBS) for 15 min. Afterward the sections were sequentially incubated with 1% Triton three times for 15 min each, 10% FBS for 30 min and with antiphospho-GluR1 (Ser-831) for 2 days at 4°C (1:200; Upstate, Barcelona, Spain). After 30 min at room temperature, sections were washed again three times in 1% Triton for 15 min each, incubated for 1 h with anti-rabbit Alexa 488 (1:200; Molecular Probes, Leiden, The Netherlands), and stained with Hoescht 33342 (Molecular Probes) for 5 min. Finally, sections were washed (TBS), mounted with DakoCytomation fluorescent mounting medium (Dako, Glostrup, Denmark), coverslipped, and examined in a Fluorescence Microscope (Zeiss Axioskop 2 Plus).

### NPY stimulated [<sup>35</sup>S]GTP $\gamma$ S binding autoradiography

The brains obtained from control and epileptic rats were cut in coronal sections (20  $\mu$ m) at -20°C, thaw mounted onto Superfrost slides, and dried on a hot plate. Tissue sections were stored at -80°C until further processing.

A slightly modified method of Sim and collaborators (44) was used. Briefly, sections were air dried for 30 min at room temperature (RT) before being rehydrated in assay buffer A (50 mM Tris-HCl, 3 mM MgCl<sub>2</sub>, 0.2 mM EGTA, 100 mM NaCl; pH 7.4) for 10 min (RT). Subsequently, a 20 min (RT) preincubation was performed in assay buffer B (assay buffer A+0.2 mM DTT, 2 mM GDP, 1  $\mu$ M DPCPX (1,3-dipropyl-8-cyclopentylxanthine), 0.5% BSA (and NPY receptor antagonist, if applicable), followed by incubation in assay buffer B + 40 pM [<sup>35</sup>S]GTP $\gamma$ S (1250 Ci/mmol, Perkin-Elmer, Denmark), 3  $\mu$ M NPY, and/or NPY receptor antagonists or PKC activator/inhibitor (Sigma Chemical Co.) for 2 h at 25°C. In each experiment, basal binding was determined by omitting the application of NPY receptor ligands and nonspecific binding by applying 10  $\mu$ M unlabeled GTP $\gamma$ S (Perkin-Elmer, Norwalk, CT, USA). The incubation was terminated by 2  $\times$  5 min washing in ice-cold 50 mM Tris-HCl buffer (pH 7.4),

followed by a brief rinse in cold deionized water. Sections were subsequently dried and exposed to Kodak BioMax MR film for 4 days together with <sup>14</sup>C standards (Amersham Life Sciences, Piscataway, NJ, USA). The films were developed in Kodak D19 developer. Optical densities were measured bilaterally over the dorsal CA1 (pyramidal layer, strata oriens, and radiatum) and CA3 (pyramidal layers, strata oriens, radiatum, and lucidum) as well as DG (molecular layer) using computer-assisted image analysis (Scion Image<sup>®</sup> analysis program). Right and left side values were averaged per section and used to calculate the mean of each animal. Background measurements immediately adjacent to each brain section were subtracted from each measurement before calculations. The percent of stimulation above basal level was calculated as [(Stim.-Basal)/Basal]\*100%.

The following compounds were used for these experiments: NPY (3  $\mu$ M, Bachem AG, Bubendorf, Switzerland), BIBP3226 (30  $\mu$ M; Bachem AG), BIIE0246 (30  $\mu$ M; Tocris Cookson Ltd., UK), L-152,804 (30  $\mu$ M; Tocris) PMA (300 nM), and BIS (3  $\mu$ M). The NPY Y<sub>2</sub> receptor population was determined using NPY in the presence of NPYY<sub>1</sub> (BIBP3226) and NPY Y<sub>5</sub> (L-152,804) receptor antagonists. Total blocking of the NPY stimulation was obtained adding a mixture of BIBP3226, BIIE246, and L-152,804.

### Statistical analysis

The data are expressed as means  $\pm$  SE. Statistics were performed using an ANOVA (one-way ANOVA), followed by Dunnett's or Bonferroni's post-tests, as indicated in the figure legends.


## RESULTS

### Modulation of glutamate release by NPY receptors in rat hippocampal synaptosomes: cross-talk with PKA or PKC pathway

We have shown that the activation of Y<sub>1</sub>, Y<sub>2</sub>, or Y<sub>5</sub> receptors with selective agonists inhibited KCl-evoked glutamate release in hippocampal synaptosomal preparations and that these inhibitory effects were prevented by their selective NPY receptor antagonists (38, 45). Moreover, we also showed that the inhibition induced by NPY (endogenous agonist, full sequence) was due to the activation of Y<sub>2</sub> receptors and not Y<sub>1</sub> or Y<sub>5</sub> subtypes (45). With the present study, we intended to clarify the possible involvement of PKA and/or PKC in the inhibitory effect mediated by NPY receptors under control and epileptic conditions.

In hippocampal synaptosomes prepared from control rats, the total KCl-evoked release of glutamate was 0.9  $\pm$  0.04 nmol glutamate/mg protein/min, and the inhibitory effect mediated by NPY (**Fig. 1A, B**) or by Y<sub>1</sub>, Y<sub>2</sub>, Y<sub>5</sub> receptor agonists (**Fig. 2A-C**) was not significantly modified by either activation of PKA with the cAMP analog 8-bromo-cAMP (500  $\mu$ M 8-Br-cAMP) or by inhibition with H89 (1  $\mu$ M). The inhibition of glutamate release induced by NPY or by the activation of Y<sub>1</sub>, Y<sub>2</sub>, Y<sub>5</sub> receptor plus 500  $\mu$ M 8-Br-cAMP was 61.5  $\pm$  6.6%, 63.0  $\pm$  3.6%, 64.7  $\pm$  3.5%, or 68.6  $\pm$  4.0% of control, respectively, and the PKA activator by itself was without effect (91.5  $\pm$  4.9% of control). Similar results were





**Figure 1.** Representative recording of the effect of NPY (1  $\mu$ M) or PMA (300 nM) (A) on the 15 mM KCl-evoked glutamate release in hippocampal synaptosomes (1 mM  $\text{Ca}^{2+}$  present in the external medium). Quantitative analysis of the effect of NPY (1  $\mu$ M) (B) on glutamate release from hippocampal synaptosomes depolarized with 15 mM KCl in the absence or presence of PKA activator (500  $\mu$ M 8-bromo-cAMP)/inhibitor (1  $\mu$ M H89) or PKC activator (300 nM PMA)/inhibitor (1  $\mu$ M BIS). Results represent the mean  $\pm$  SE of 4 to 9 independent experiments in different synaptosomal preparations.  $**P < 0.01$ , Dunnett's post-test, statistical significance when compared with control (KCl stimulation).


obtained in the presence of 1  $\mu$ M NPY (Fig. 1B), 1  $\mu$ M [Leu<sup>31</sup>, Pro<sup>34</sup>]NPY (Fig. 2A), 300 nM NPY(13–36) (Fig. 2B), or 1  $\mu$ M NPY(19–23)-(Gly<sup>1</sup>, Ser<sup>3</sup>, Gln<sup>4</sup>, Thr<sup>6</sup>, AL<sup>31</sup>, Aib<sup>32</sup>, Gln<sup>34</sup>)-PP (Fig. 2C) plus 1  $\mu$ M H89 as follows: 65.8  $\pm$  3.3%, 59.3  $\pm$  3.1%, 63.8  $\pm$  1.8%, 70.4  $\pm$  3.5% of control, respectively. Again, the inhibitor of PKA (H89) by itself had no effect (91.8  $\pm$  4.3% of control). All these observations allow us to conclude that PKA was not interfering with the inhibition of glutamate release mediated by NPY receptor activation.

PKC-mediated substrate phosphorylation has been implicated in a broad spectrum of neuronal events, including neurotransmitter release (28). Indeed, activation of PKC by phorbol esters leads to an increase in neurotransmitter release from nerve terminals of hippocampal slices (46), from nerve terminals in the spinal


cord (47), and from isolated nerve terminals from mammalian brain (48, 49). In agreement with these studies, the activation of PKC with 300 nM PMA induced an increase in KCl-evoked glutamate release to 129.7  $\pm$  3.0% of control (Figs. 1, 2), which was blocked by the PKC inhibitor (BIS) to 99.3  $\pm$  6.6% of control (Figs. 1B, 2). By itself, 1  $\mu$ M BIS did not induce any significant effect (92.5  $\pm$  5.4% of control). Moreover, the inhibitory effect induced by 1  $\mu$ M [Leu<sup>31</sup>, Pro<sup>34</sup>]NPY (Fig. 2A) or 1  $\mu$ M NPY(19–23)-(Gly<sup>1</sup>, Ser<sup>3</sup>, Gln<sup>4</sup>, Thr<sup>6</sup>, AL<sup>31</sup>, Aib<sup>32</sup>, Gln<sup>34</sup>)-PP (Fig. 2C) was not affected by the simultaneous activation of PKC (100.0  $\pm$  7.3% or 93.8  $\pm$  8.0% of control, respectively), keeping in mind that PMA by itself increased glutamate release to 129.7  $\pm$  3.0% of control. However, this was not the case concerning NPY or Y<sub>2</sub> receptor activation, since in the simultaneous presence of NPY or Y<sub>2</sub> receptor agonist plus PMA, glutamate release was 133.2  $\pm$  2.2% (Fig. 1B) or 133.8  $\pm$  2.9% (Fig. 2B) of control, respectively. Moreover, the inhibitory effect mediated by 1  $\mu$ M NPY (66.0  $\pm$  4.7% of control) or 300 nM NPY(13–36) (71.0  $\pm$  1.2% of control) was recovered in the presence of the PKC inhibitor to 68.9  $\pm$  2.6% or 65.0  $\pm$  5.4% of control, respectively (Fig. 1B, Fig. 2B). The PKC inhibitor by itself (1  $\mu$ M BIS) did not affect the inhibition induced by Y<sub>1</sub> (Fig. 2A), Y<sub>2</sub> (Fig. 2B), Y<sub>5</sub> (Fig. 2C) receptor agonists or NPY (Fig. 1B) (75.8  $\pm$  2.9%, 67.0  $\pm$  5.5%, 63.6  $\pm$  3.4%, or 68.2  $\pm$  3.2% of control, respectively). These results show that, under control conditions, the pharmacological activation of PKC blocks the inhibition of glutamate release mediated by the activation of Y<sub>2</sub> receptors.

### Involvement of PKC in the modulation of glutamate release by Y<sub>2</sub> receptors in hippocampal synaptosomes obtained from epileptic rats

Several studies have demonstrated significant alterations in PKC expression and subcellular distribution in the hippocampus after KA treatment (50, 51). To investigate the role of PKC in the inhibition of glutamate release induced by Y<sub>2</sub> receptor activation under epileptic conditions, we injected rats with 10 mg/kg KA (i.p.), a well-established model of epileptogenesis, as described in Materials and Methods. Using synaptosomes obtained from rats sacrificed 6 h postinjection, we observed that the total KCl-evoked release of glutamate was 1.1  $\pm$  0.03 nmol glutamate/mg protein/min and that none of the agonists alone or in the presence of the PKC activator had an inhibitory effect (Fig. 3A). The values obtained with 1  $\mu$ M NPY, 300 nM NPY(13–36), 1  $\mu$ M NPY plus 300 nM PMA, or 300 nM NPY(13–36) plus 300 nM PMA were 94.1  $\pm$  3.3%, 93.4  $\pm$  3.3%, 93.7  $\pm$  3.1%, or 96.0  $\pm$  4.5% of control, respectively (Fig. 3A). Moreover, 1  $\mu$ M BIS did not modify the KCl-evoked glutamate release (108.0  $\pm$  4.0% of control); in contrast to what happened under control conditions, 300 nM PMA by itself had no effect (100.2  $\pm$  3.9% of control) (Fig. 3A). However, after inhibition of PKC




**Figure 2.** Involvement of PKA or PKC on the inhibition of glutamate release mediated by NPY Y<sub>1</sub>, Y<sub>2</sub>, or Y<sub>5</sub> receptor activation in rat hippocampal synaptosomes. Quantitative analysis of the effect of 1 μM [Leu<sup>31</sup>, Pro<sup>34</sup>]NPY (A), 300 nM NPY(13–36) (B), 1 μM NPY(19–23)-(Gly<sup>1</sup>, Ser<sup>3</sup>, Gln<sup>4</sup>,



**Figure 3.** Involvement of PKC on the inhibition of glutamate release mediated by NPY or Y<sub>2</sub> receptor activation in epileptic rats. Quantitative analysis of the effect of 1 μM NPY or 300 nM NPY(13–36) on glutamate release evoked by 15 mM KCl depolarization in hippocampal synaptosomes obtained from epileptic rats sacrificed 6 h (A) or 24 h (B) postinjection. Status epilepticus was induced in rats after injection with KA (10 mg/kg) as described in Material and Methods. Results represent the mean ± SE of 3 to 7 independent experiments in different synaptosomal preparations. \*\**P* < 0.01, Dunnett's post-test, statistical significance when compared with control (KCl stimulation).

Thr<sup>6</sup>, AL<sup>31</sup>, Aib<sup>32</sup>, Gln<sup>34</sup>)-PP (C) on glutamate release from hippocampal synaptosomes depolarized with 15 mM KCl in the absence or in the presence of PKA activator (500 μM 8-bromo-cAMP)/inhibitor (1 μM H89) or PKC activator (300 nM PMA)/inhibitor (1 μM BIS). Results represent the mean ± SE of 4 to 9 independent experiments in different synaptosomal preparations. \*\**P* < 0.01, Dunnett's post-test, statistical significance when compared with control (KCl stimulation).




**Figure 4.** Western blot analysis of the changes in GluR1 (A, B) or phospho-GluR1 (C, D) levels in hippocampal subregions (CA1, CA3, DG) of control rats (injected with saline) and epileptic rats sacrificed 6 h (A, C) or 24 h (B, D) after KA injection (10 mg/kg, i.p.). Results are expressed as mean percentage of control  $\pm$  SE of 3 or 4 independent experiments. \*\*\* $P$  < 0.001, \*\* $P$  < 0.01. Bonferroni's post-test, statistical significance when compared with CA1, CA3, or DG from control rats.

with 1  $\mu$ M BIS, we again observed a significant inhibitory effect caused by 1  $\mu$ M NPY (70.0  $\pm$  3.6% of control) or by 300 nM NPY(13–36) (71.0  $\pm$  2.3% of control) (Fig. 3A). Taking in consideration that 6 h after kainate administration cells can be in a depolarized state and still have increased calcium levels, we performed additional experiments in which we observed that in synaptosomes obtained from epileptic rats sacrificed 6 h postinjection, the P-/Q-type VGCC blocker  $\omega$ -agatoxin IVA (100 nM) inhibited glutamate release to 51% of control (data not shown), similar to what we observed earlier in control animals (45). These results show that the functional release of glutamate is fully functional in epileptic rats and is not blocked due to increased levels of presynaptic calcium.

The results obtained 24 h post-injection were similar to the control situation (Fig. 3B). The total KCl-evoked release of glutamate was 1.1  $\pm$  0.1 nmol glutamate/mg protein/min; 1  $\mu$ M NPY or 300 nM NPY(13–36) inhibited KCl-evoked glutamate release to 62.0  $\pm$  1.4% or 68.6  $\pm$  1.4% of control, respectively (Fig. 3B). Again, PMA by itself potentiated the release of glutamate to 136.3  $\pm$  3.1% of control, and in the presence of NPY or the Y<sub>2</sub> receptor agonist, glutamate release was 137.8  $\pm$  4.0% or 135.0  $\pm$  3.7% of control, respectively. Moreover, as observed un-

der control conditions, 1  $\mu$ M BIS by itself was without effect (98.2  $\pm$  7.6% of control) and completely blocked the potentiation of glutamate release induced by PMA (105.0  $\pm$  7.0% of control) (Fig. 3B). Based on these results, we suggest that at 6 h postinjection PKC is highly active and occludes the inhibition of glutamate release mediated by Y<sub>2</sub> receptors.



**Figure 5.** Representative fluorescence microscopy images of phospho-GluR1 (Ser-831) immunoreactivity in the CA1 pyramidal cell layer of the hippocampus of control rats (A) or epileptic rats injected with KA (10 mg/kg) and sacrificed 6 h (B) or 24 h (C) postinjection. There is an increase of immunoreactivity against phospho-GluR1 (Ser-831) in animals with seizures. At 6 h postinjection (B), punctuate-like labeling is clearly visible, suggesting a synaptic localization (arrows), whereas after 24 h (C) immunoreactivity is distributed along neurites (arrows). Brain slices (20  $\mu$ m) were labeled with antiphospho-GluR1 (Ser-831) (green) and stained with Hoescht 33342 (blue). Scale: 50  $\mu$ m.



TABLE 1. Effect of PKC activator (PMA) or inhibitor (BIS) on NPY-stimulated [<sup>35</sup>S]GTPγS binding in adult control and epileptic rat hippocampal subregions<sup>a</sup>

Region	Percentage increase above basal [ <sup>35</sup> S]GTPγS binding (mean±SE, n = 6–8)						
	NPY			Y <sub>2</sub> <sup>b</sup>			
	+PMA	+BIS	+PMA+BIS	+PMA	+PMA+BIS		
<i>CA1</i>							
Control	63 ± 6	51 ± 6	66 ± 7	66 ± 11	33 ± 3	31 ± 9	45 ± 5
Epileptic	82 ± 9	58 ± 13	74 ± 14	78 ± 18	41 ± 5	38 ± 18	59 ± 10
<i>CA3</i>							
Control	70 ± 12	57 ± 9	75 ± 11	74 ± 12	56 ± 10	54 ± 13	58 ± 13
Epileptic	74 ± 11	59 ± 9	72 ± 12	74 ± 15	63 ± 12	48 ± 10	73 ± 13
<i>DG</i>							
Control	41 ± 5	33 ± 5	45 ± 7	48 ± 9	0	0	2 ± 5
Epileptic	57 ± 8	45 ± 8	52 ± 9	57 ± 9	10 ± 9	1 ± 13	18 ± 7

<sup>a</sup>Control rats were injected with saline (0.9% NaCl); epileptic seizures were induced with KA (10 mg/kg, i.p. injection) and rats were sacrificed 6 h postinjection. Brain sections were processed as described in Materials and Methods. CA1, regio-superior cornu ammonis; CA3, regio-inferior cornu ammonis; DG, dentate gyrus. <sup>b</sup>The NPY Y<sub>2</sub> receptor population was determined using NPY in the presence of NPY Y<sub>1</sub> (BIBP3226) and NPY Y<sub>5</sub> (L-152,804) receptor antagonists.

### GluR1 and phospho-GluR1 (Ser-831) levels in hippocampal subregions of control and epileptic rats

The results described above suggest that in the acute phase of epilepsy, PKC may be strongly activated and that this effect can be associated with the loss of Y<sub>2</sub> receptor-mediated inhibition of glutamate release. To assess whether PKC is active under these conditions, we decided to investigate the levels of phospho-GluR1 (Ser-831) as an indication of PKC activity. AMPA receptors containing the GluR1 subunit may be regulated by extracellular signals acting through PKC (52, 53). Indeed, PKC phosphorylates Ser-831 of the GluR1 subunit (54) and can contribute to modulate synaptic transmission. Thus, we evaluated the levels of GluR1 and phospho-GluR1 in hippocampal subregions of control and epileptic rats.

In hippocampal subregions CA1, CA3, and DG, GluR1 levels at 6 h or 24 h postinjection were not significantly different from those of control (Fig. 4A, B). In contrast, phospho-GluR1 (Ser-831) levels robustly increased at 6 h (253.7±25.3% of control), and even more pronounced 24 h postinjection in the CA1 subregion (1325.6±267.4% of control) (Fig. 4C, D). No significant effects were seen in the DG at any time point after KA injection (Fig. 4C, D).


In the CA1 subregion, we observed an increase of immunoreactivity against phospho-GluR1 (Ser-831) in animals with seizures (Fig. 5). At 6 h postinjection (Fig. 5B), punctuate-like labeling was clearly visible, suggesting a synaptic localization, whereas after 24 h the immunoreactivity was distributed along neurites (Fig. 5C).

### Characterization of the intracellular mechanisms by which PKC activity blocks NPY-mediated inhibition of glutamate release

To study both the distribution and functionality of the NPY receptors under control and epileptic conditions,

we performed functional autoradiographic studies. Table 1 and Fig. 6A–C show levels of [<sup>35</sup>S]GTPγS binding in CA1, CA3, and DG after stimulation with 3 μM NPY. This effect was completely blocked by a mixture of Y<sub>1</sub>, Y<sub>2</sub>, and Y<sub>5</sub> receptor antagonists (30 μM BIBP3226+30 μM BIIE0246+30 μM L-152,804) (Fig. 6D). Activation of the Y<sub>2</sub> receptor subtype (3 μM NPY+30 μM BIBP3226+30 μM L-152,804) only induced an increased binding signal above basal binding in the CA1 and CA3 subregions, correlating with the binding pattern observed for the Y<sub>2</sub> receptor in traditional autoradiographic studies (55) (Table 1; Fig. 6E). Moreover, the PKC activator (PMA) and/or inhibitor (BIS) did not significantly alter the total NPY receptor or Y<sub>2</sub> receptor-mediated [<sup>35</sup>S]GTPγS binding under either control or epileptic conditions (6 h postinjection) (Table 1; Fig. 6). These results suggest that the function of the NPY receptors is similar under control *vs.* epileptic conditions. Moreover, PKC does not seem to directly modulate the activity of NPY receptors, suggesting that the PKC target is located downstream the receptor level.

We previously showed that none of the NPY receptor agonists used currently could reduce ionomycin-induced glutamate release (45). Accordingly, in Fig. 7 we observed that PMA by itself (100.0±3.9% of control) or in the presence of NPY (86.5±2.3% of control) or NPY(13–36) (93.8±2.5% of control) did not significantly potentiate or inhibit glutamate release, in contrast to what happened when we stimulated glutamate release with KCl (Fig. 1B, Fig. 2B); 1 μM BIS or 1 μM BIS + 1 μM NPY had no effect (data not shown). Since ionomycin promotes calcium influx and glutamate release independently of voltage-gated Ca<sup>2+</sup> channels (VGCCs) activity, the present results suggest that PKC is acting on VGCCs, increasing Ca<sup>2+</sup> conductance and thereby occluding the inhibitory effect of Y<sub>2</sub> receptors.




**Figure 6.** Representative images of [<sup>35</sup>S]GTPγS functional binding in the rat hippocampal CA1, CA3, and DG at 6 h after i.p. injection of saline (0.9% NaCl) or KA (10 mg/kg). Visualization of all NPY receptor subtypes (Y<sub>1</sub>, Y<sub>2</sub>, Y<sub>5</sub>) with 3 μM NPY-stimulated binding (A), nonspecific binding (B), basal binding (C), and total block of NPY-stimulated binding (3 μM NPY+30 μM Y<sub>1</sub>, Y<sub>2</sub>, Y<sub>5</sub> antagonists) (D) in saline-treated rats. Y<sub>2</sub> receptor functional binding (3 μM NPY+30 μM Y<sub>1</sub>, Y<sub>5</sub> antagonists) did not differ in CA1 and CA3 (arrows) between saline- (E) and KA-treated rats (F). No functional Y<sub>2</sub> receptor binding was detected in the DG. The presence of PKC activator (300 nM PMA) did not affect Y<sub>2</sub> receptor-mediated functional binding in either saline- (G) or KA-injected rats (H).

## DISCUSSION

In previous studies, we and others have shown that NPY receptors modulate Ca<sup>2+</sup> influx (38, 56), K<sup>+</sup> currents (57, 58), and glutamate release (38, 59) from neuronal cells. Moreover, NPY inhibits excitatory neurotransmission in the hippocampus (18), exogenous administration of NPY prominently suppresses limbic seizure activity induced by KA (15), and NPY-deficient mice develop more severe seizures in response to KA, suggesting that NPY may act as an endogenous anticonvulsant agent. So, since NPY affects glutamatergic synaptic

transmission and neuronal excitability (5, 10, 12), it is of great interest to investigate the functional role of the NPY system in epileptic seizures. Indeed, little is known about the signaling mechanism (mechanisms) that underlie NPY receptor effects under epileptic conditions.

In common with some other presynaptic metabotropic receptors (60), the observed inhibitory effect mediated by NPY Y<sub>1</sub>, Y<sub>2</sub>, and Y<sub>5</sub> receptors did not work through a cyclic AMP-dependent mechanism as it was not influenced by PKA activator (8-Br-cAMP) or by its inhibitor (H89). In fact, in our study we observed that the activation or inhibition of PKA failed to modify inhibition of glutamate release mediated by any of the NPY receptors. In contrast, the role of PKC activation in regulation of presynaptic function in general, and glutamate release in particular, is well established in synaptosomal studies (29, 60). In the present work, we observed that the activation of PKC by a phorbol ester (PMA) induced an increase of glutamate release. In fact, it is well established that the activation of PKC by phorbol esters leads to an increase in neurotransmitter release from rat hippocampal slices (46), and mammalian brains synaptosomes (29, 48), and also potentiates excitatory postsynaptic currents (EPSCs) (61). NPY- or Y<sub>2</sub> receptor-mediated inhibition of glutamate release under control conditions was abolished following PKC activation. In contrast, when using synaptosomes obtained from epileptic rats 6 h postinjection, we observed no NPY receptor-mediated inhibition of glutamate release (21). Moreover, phorbol esters did not affect glutamate release, in contrast to what was observed 24 h postinjection and in controls. These results suggest that PKC can be highly active following status epilepticus and somehow blocks the inhibition of glu-



**Figure 7.** Lack of effect of NPY, NPY(13–36), and PMA on glutamate release evoked by 5 μM ionomycin in control rat hippocampal synaptosomes. Results represent the mean ± SE of 4 to 8 independent experiments in different synaptosomal preparations.



tamate release induced by  $Y_2$  receptor activation. Another possible interpretation for our results is the inhibition of PKC effects by NPY acting through  $Y_1$  and  $Y_5$  receptors. As we show in the present study, in epileptic rats (6 h post-kainate injection), NPY is not efficient at inhibiting the release of glutamate; this effect is similar to that observed after stimulation of  $Y_2$  receptors but different from the results obtained in the presence of  $Y_1$  or  $Y_5$  agonists (21). The effect of NPY under basal conditions and epileptic ones (24 h postinjection) is mediated mainly through  $Y_2$  receptors, but after status epilepticus (6 h), when PKC is active,  $Y_2$  receptor activity is not linked to the efficient inhibition of glutamate release. However, whether or not  $Y_1$  and  $Y_5$  receptor activity can shortcut the effect of PKC in stimulating the release of glutamate is an alternative and attractive hypothesis. Future studies will determine whether in epileptic rats (6 h) the blockade of  $Y_2$  receptors relieve the predominant effect of NPY at  $Y_2$  receptors, making  $Y_1$  and  $Y_5$  receptors sensitive to NPY (as we previously reported in 45), and whether in this condition NPY can efficiently block the stimulatory effect of PKC.

It is clear that status epilepticus induces alterations in the subcellular distribution of individual PKC isoforms in a temporally and regionally specific manner (62). In fact, we observed that by inhibiting PKC in synaptosomes isolated 6 h postinjection with KA,  $Y_2$  receptors recovered their inhibitory effect on glutamate release. Moreover, analyzing the expression levels of phospho-GluR1 (Ser-831), it is evident there is an increase of PKC activity after status epilepticus. This increase was significant in the CA1 subregion, in agreement with other studies that show a differential activity of PKC among brain regions or subregions (62). Also, Hussain and Carpenter (63) showed that the presence and function of PKC are age dependent and different between Schaffer collateral-CA1 and Mossy fibers-CA3 pathways within the normal hippocampus. With the present data, we cannot form definitive conclusions about the mechanisms contributing to the differences observed in the functional release of glutamate between 6 h and 24 h. We can, however, suggest that other mechanisms able to shortcut/silent the stimulatory effect of PKC on glutamate release (*i.e.*, increased activity of phosphatases) may be functional at 24 h but not at 6 h postinjection. It is important to keep in mind that phospho-GluR1 was used as a marker PKC activity, but this approach cannot lead to conclusions about the spatial and temporal activity of PKC at presynaptic sites, especially important for the control of voltage-gated calcium channels.

As already mentioned, several actions have been attributed to PKC, including not only the enhancement of  $Ca^{2+}$  currents and elevation of cytosolic-free  $Ca^{2+}$  resulting in increased neurotransmitter release (64), inhibition of  $K^+$  channels (31), and activation of exocytotic machinery downstream of  $Ca^{2+}$  influx (30), but also regulation of activity of NMDA receptors (65) or

$\alpha_{1d}$ -adrenergic receptors (66). However, the mechanism (or mechanisms) whereby PKC blocks the inhibitory effect of  $Y_2$  receptors is unknown. Therefore, we first tried to investigate whether PKC was acting directly on the receptors or at the G-protein level. Our results indicated that this might not be the case, in contrast to what is known to happen with several other receptors (65). Indeed, the [ $^{35}S$ ]GTP $\gamma$ S binding assay we used to assess the function of  $Y_2$  receptors gave similar results under control and epileptic conditions with or without the pharmacological activation of PKC with phorbol ester, indicating that these receptors are not functionally directly regulated by PKC. So the precise mechanism (or mechanisms) underlying the blockage of  $Y_2$  receptors are yet to be determined, but given that modulation of ion channel activity is considered one of the major target mechanisms, it is plausible that a good target candidate for PKC could be the voltage-gated  $Ca^{2+}$  channels (VGCCs). In previous studies performed by us and others, it was shown that NPY receptors modulate (inhibit) the [ $Ca^{2+}$ ] $_i$  response in the hippocampus, and this was mainly due to the inhibition of different VGCCs (45). If PKC is acting on VGCCs by increasing  $Ca^{2+}$  conductance, as we suggested in the present study and as described by others (29), it is plausible to suggest that this mechanism may account for the cross-talk between PKC and  $Y_1$  (29) or PKC and  $Y_2$  (present study) receptor function. A major unbalance of the excitatory/inhibitory transmission occurs in status epilepticus and causes a major shift toward hyperexcitability, contributing to the generalization of seizure activity and to tonic-clonic convulsions. Our present demonstration that PKC contributes to the increased release of glutamate and at the same time, blocks a key inhibitory modulator ( $Y_2$  receptor) may help to highlight the impact of the PKC/NPY cross-talk in status epilepticus.

In summary, the present findings show a transient loss of NPY- and  $Y_2$  receptor-mediated inhibition of glutamate release following status epilepticus. We also show that PKC is highly active and may contribute to occlude the functional effects of  $Y_2$  receptors in inhibiting glutamate release. Since  $Y_2$  receptors are essential for NPY regulation of limbic seizures originating in the hippocampus, understanding the cell signaling that underlies its effect under control and epileptic conditions may be quite useful to find potential new treatments for temporal lobe epilepsy. EJ

We acknowledge Dr. Henri Doods (Boehringer Ingelheim Pharma KG) for generously providing us the BIIE0246 compound, and Dr. Inês Araújo for providing hippocampal lysates of 24 h epileptic rats. This work was supported by Fundação para a Ciência e a Tecnologia, Portugal (Project POCTI/NSE/46848/2002, POCTI/FCB/46804/2002 and Grant BPD/11484/2002), FEDER, the Danish National Research Council project No. 64750, the Friis Foundation, the Lundbeck Foundation, and a grant from Fundação Calouste Gulbenkian, Portugal.

## REFERENCES

- Wettstein, J. G., Earley, B., and Junien, J. L. (1995) Central nervous system pharmacology of neuropeptide Y. *Pharmacol. Ther.* **65**, 397–414
- Berglund, M. M., Hipskind, P. A., and Gehlert, D. R. (2003) Recent developments in our understanding of the physiological role of PP-fold peptide receptor subtypes. *Exp. Biol. Med.* **228**, 217–244
- Silva, A. P., Xapelli, S., Grouzmann, E., and Cavadas, C. (2005) The putative neuroprotective role of neuropeptide Y in the central nervous system. *Curr. Drug. Targets. CNS. Neurol. Disord.* **4**, 331–347
- Wahlestedt, C., and Reis, D. J. (1993) Neuropeptide Y-related peptides and their receptors—are the receptors potential therapeutic drug targets? (Review) *Ann. Rev. Pharmacol. Toxicol.* **32**, 309–352
- Vezzani, A., Sperk, G., and Colmers, F. (1999) Neuropeptide Y: emerging evidence for a functional role in seizure modulation. *Trends Neurosci.* **22**, 25–30
- Colmers, W. F., and El Bahh, B. (2003) Neuropeptide Y and epilepsy. *Epilepsy Curr.* **3**, 53–58
- Woldbye, D. P., and Kokaia, M. (2004) Neuropeptide Y and seizures: effects of exogenously applied ligands. *Neuropeptides* **38**, 253–260
- Michel, M. C., Beck-Sickinger, A., Cox, H., Doods, H. N., Herzog, H., Larhammar, D., Quirion, R., Schwartz, T., and Westfall, T. (1998) International union of pharmacology recommendations for the nomenclature of neuropeptide Y, peptide YY, and pancreatic polypeptide receptors. *Pharmacol. Rev.* **50**, 143–150
- Parker, R. M. C., and Herzog, H. (1999) Regional distribution of Y-receptor subtype mRNA in rat brain. *Eur. J. Neurosci.* **11**, 1431–1448
- Wu, Y. F., and Li, S. B. (2005) Neuropeptide Y expression in mouse hippocampus and its role in neuronal excitotoxicity. *Acta Pharmacol. Sin.* **26**, 63–68
- Gobbi, M., Gariboldi, M., Piwko, C., Hoyer, D., Sperk, G., and Vezzani, A. (1998) Distinct changes in peptide YY binding to, and mRNA levels of, Y1 and Y2 receptors in the rat hippocampus associated with kindling epileptogenesis. *J. Neurochem.* **70**, 1615–1622
- El Bahh, B., Balosso, S., Hamilton, T., Herzog, H., Beck-Sickinger, A. G., Sperk, G., Gehlert, D. R., Vezzani, A., Colmers, W. F. (2005) The anti-epileptic actions of neuropeptide Y in the hippocampus are mediated by Y<sub>1</sub> and not Y<sub>5</sub> receptors. *Eur. J. Neurosci.* **22**, 1417–1430
- Woldbye, D. P., Nanobashvili, A., Sorensen, A. T., Husum, H., Bolwig, T. G., Sorensen, G., Ernfors, P., and Kokaia, M. (2005) Differential suppression of seizures via Y<sub>2</sub> and Y<sub>5</sub> neuropeptide Y receptors. *Neurobiol. Dis.* **20**, 760–772
- Erickson, J. C., Clegg, K. E., and Palmiter, R. D. (1996) Sensitivity to leptin and susceptibility to seizures of mice lacking neuropeptide Y. *Nature* **381**, 415–421
- Baraban, S. C., Hollopeter, G., Erickson, J. C., Schwartzkroin, P. A., and Palmiter, R. D. (1997) Knock-out mice reveal a critical antiepileptic role for neuropeptide Y. *J. Neurosci.* **17**, 8927–8936
- Vezzani, A., Michalkiewicz, M., Michalkiewicz, T., Moneta, D., Ravizza, T., Richichi, C., Aliprandi, M., Mulé, F., Gobbi, M., Schwarzer, C., and Sperk, G. (2002) Seizure susceptibility and epileptogenesis are decreased in transgenic rats overexpressing neuropeptide Y. *Neuroscience* **110**, 237–243
- Richichi, C., Lin, E. J., Stefanin, D., Colella, D., Ravizza, T., Grignaschi, G., Veglianesi, P., Sperk, G., During, M. J., and Vezzani, A. (2004) Anticonvulsant and antiepileptogenic effects mediated by adeno-associated virus vector neuropeptide Y expression in the rat hippocampus. *J. Neurosci.* **24**, 3051–3059
- Colmers, W. F., Klapstein, G. J., Fournier, A., St-Pierre, S., and Treherne, K. A. (1991) Presynaptic inhibition by neuropeptide Y in rat hippocampal slice *in vitro* is mediated by a Y<sub>2</sub> receptor. *Br. J. Pharmacol.* **102**, 41–44
- Bleakman, D., Harrison, N. L., Colmers, W. F., and Miller, R. J. (1992) Investigations into neuropeptide Y-mediated presynaptic inhibition in cultured hippocampal neurones of the rat. *Br. J. Pharmacol.* **107**, 334–340
- Greber, S., Schwarze, C., and Sperk, G. (1994) Neuropeptide Y inhibits potassium-stimulated glutamate release through Y<sub>2</sub> receptors in rat hippocampal slices *in vitro*. *Br. J. Pharmacol.* **113**, 737–740
- Silva, A. P., Xapelli, S., Pinheiro, P. S., Ferreira, R., Lourenço, J., Cristovão, A., Grouzmann, E., Cavadas, C., Oliveira, C. R., and Malva, J. O. (2005) Up-regulation of neuropeptide Y levels and modulation of glutamate release through neuropeptide Y receptors in the hippocampus of kainate-induced epileptic rats. *J. Neurochem.* **93**, 163–170
- Klapstein, G. J., and Colmers, W. F. (1997) Neuropeptide Y suppresses epileptiform activity in rat hippocampus *in vitro*. *J. Neurophysiol.* **78**, 1651–1661
- Woldbye, D. P. D., Larsen, P. J., Mikkelsen, J. D., Klemp, K., Madsen, T. M., and Bolwig, T. G. (1997) Powerful inhibition of kainic acid seizures by neuropeptide Y via Y<sub>5</sub>-like receptors. *Nature Med.* **3**, 761–764
- Cambrey-Deakin, M. A., Adu, J., and Burgoyne, R. D. (1990) Neurogenesis in cerebellar granule cells *in vitro*: a role for protein kinase C. *Brain Res. Dev. Brain Res.* **53**, 40–46
- Campanot, R. B., Draker, D. D., and Senger, D. L. (1994) Evidence that protein kinase C activities involved in regulating neurite growth are localized to distal neurites. *J. Neurochem.* **63**, 868–878
- Parrow, V., Fagerstrom, S., Meyerson, G., Nanberg, E., and Pahlman, S. (1995) Protein kinase C- $\alpha$  and - $\epsilon$  are enriched in growth cones of differentiating SH-SY5Y human neuroblastoma cells. *J. Neurosci. Res.* **41**, 782–791
- Felipo, V., Minana, M. D., and Grisolia, S. (1993) Inhibitors of protein kinase C prevent the toxicity of glutamate in primary neuronal cultures. *Brain Res.* **604**, 192–196
- Robinson, P. J. (1992) The role of protein kinase C and its neuronal substrates dephosphin, B-50, and MARCKS in neurotransmitter release. *Mol. Neurobiol.* **5**, 87–130
- Wang, S. J. (2006) Facilitatory effect of aspirin on glutamate release from rat hippocampal nerve terminals: involvement of protein kinase C pathway. *Neurochem. Int.* **48**, 181–190
- Redman, R. S., Searl, T. J., Hirsh, J. K., Silinsky, E. M. (1997) Opposing effects of phorbol esters on transmitter release and calcium currents at frog motor nerve endings. *J. Physiol.* **15**, 41–48
- Hoffman, D. A., and Johnston, D. (1998) Downregulation of transient K<sup>+</sup> channels in dendrites of hippocampal CA1 pyramidal neurons by activation of PKA and PKC. *J. Neurosci.* **15**, 3521–3528
- Chow, K. B., Jones, R. L., and Wise, H. (2003) Protein kinase A-dependent coupling of mouse prostacyclin receptors to Gi is cell-type dependent. *Eur. J. Pharmacol.* **474**, 7–13
- Bijak, M. (1999) Neuropeptide Y suppresses epileptiform activity in rat frontal cortex and hippocampus *in vitro* via different NPY receptor subtypes. *Neurosci. Lett.* **268**, 115–118
- Marsh, D. J., Baraban, S. C., Hollopeter, G., and Palmiter, R. D. (1999) Role of the Y<sub>5</sub> neuropeptide Y receptor in limbic seizures. *Proc. Natl. Acad. Sci. U. S. A.* **96**, 13518–13523
- Sperk, G., Lassmann, H., Baran, H., Seitelberger, F., and Hornykiewicz, O. (1985) Kainic acid-induced seizures: dose-relationship of behavioural, neurochemical and histopathological changes. *Brain Res.* **338**, 289–295
- McMahon, T. H., Foran, P., Dolly, J. O., Verhage, M., Wiegant, V. M., and Nicholls, D. G. (1992) Tetanus toxin and botulinum toxins type A and B inhibit glutamate,  $\gamma$ -aminobutyric acid, aspartate, and Met-enkephalin release from synaptosomes. *J. Biol. Chem.* **267**, 21338–21343
- Malva, J. O., Carvalho, A. P., and Carvalho, C. M. (1996) Domoic acid induces the release of glutamate in the rat hippocampal CA3 sub-region. *NeuroReport* **7**, 1330–1344
- Silva, A. P., Carvalho, A. P., Carvalho, C. M., and Malva, J. O. (2001) Modulation of intracellular calcium changes and glutamate release by neuropeptide Y<sub>1</sub> and Y<sub>2</sub> receptors in the rat hippocampus: differential effects in CA1, CA3 and dentate gyrus. *J. Neurochem.* **79**, 286–296
- Whittaker, V. P. (1984) The synaptosome. In *Handbook of Neurochemistry* (Lajtha, A., eds) Vol. 7, pp. 1–39, Plenum Press, New York
- Layne, E. (1957) Spectrophotometric and turbidimetric methods for measuring proteins. In *Methods in Enzymology* (Colowick,

- S. P., and Kaplan, N. O., eds) Vol. 3, pp. 447–451, Academic, New York
41. Smith, P. K., Krohn, R. I., Hermanson, G. T., Mallia, A. K., Gartner, F. H., Provenzano, M. D., Fujimoto, E. K., Goeke, N. M., Olson, B. J., and Klen, D. C. (1985) Measurement of protein using bicinchoninic acid. *Anal. Biochem.* **150**, 76–85
  42. Nicholls, D. G., Sihra, T. S., and Sanchez-Prieto, J. (1987) Calcium-dependent and -independent release of glutamate from synaptosomes monitored by continuous fluorometry. *J. Neurochem.* **49**, 50–57
  43. Pinheiro, P. S., Rodrigues, R. J., Rebola, N., Xapelli, S., Oliveira, C. R., and Malva, J. O. (2005) Presynaptic kainate receptors are localized close to release sites in rat hippocampal synapses. *Neurochem. Int.* **47**, 309–316
  44. Sim, L. J., Selley, D. E., and Childers, S. R. (1995) In vitro autoradiography of receptor-activated G proteins in rat brain by agonist-stimulated guanylyl 5'-[gamma-<sup>35</sup>S]thio]-triphosphate binding. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 7242–7246
  45. Silva, A. P., Carvalho, A. P., Carvalho, C. M., and Malva, J. O. (2003) Functional interaction between neuropeptide Y receptors and modulation of calcium channels in the rat hippocampus. *Neuropharmacology* **42**, 282–292
  46. Parfitt, K. D., and Madison, D. V. (1993) Phorbol esters enhance synaptic transmission by a presynaptic, calcium-dependent mechanism in rat hippocampus. *J. Physiol.* **471**, 245–268
  47. Gandhi, V. C., and Jones, D. J. (1992) Protein kinase C modulates the release of [<sup>3</sup>H]5-hydroxytryptamine in the spinal cord of the rat: the role of L-type voltage-dependent calcium channels. *Neuropharmacology* **31**, 1101–1109
  48. Herrero, L., Miras-Portugal, M. T., and Sánchez-Prieto, J. (1992) Activation of protein kinase C by phorbol esters and arachidonic acid required for the optimal potentiation of glutamate exocytosis. *J. Neurochem.* **59**, 1574–1577
  49. Coffey, E. T., Sihra, T. S., and Nicholls, D. G. (1993) Protein kinase C and the regulation of glutamate exocytosis from cerebrocortical synaptosomes. *J. Biol. Chem.* **268**, 21060–21065
  50. Guglielmetti, F., Rattray, M., Baldessari, S., Butelli, E., Samanin, R., and Bendotti, C. (1997) Selective up-regulation of protein kinase C in granule cells after kainic acid-induced seizures in rat. *Brain Res. Mol. Brain Res.* **49**, 188–196
  51. McNamara, R. K., and Lenox, R. H. (2000) Differential regulation of primary protein kinase C substrate (MARCKS, MLP, GAP 43, RC3) mRNAs in the hippocampus during kainic acid-induced seizures and synaptic reorganization. *J. Neurosci. Res.* **62**, 416–426
  52. Tan, S. E., Wenthold, R. J., and Soderling, T. R. (1994) Phosphorylation of AMPA-type glutamate receptors by calcium/calmodulin-dependent protein kinase II and protein kinase C in cultured hippocampal neurons. *J. Neurosci.* **14**, 1123–1129
  53. Blackstone, C., Murphy, T. H., Moss, S. J., Baraban, J. M., and Huganir, R. L. (1994) Cyclic AMP and synaptic activity-dependent phosphorylation of AMPA-preferring glutamate receptors. *J. Neurosci.* **14**, 7585–7593
  54. Roche, K. W., O'Brien, R. J., Mammen, A. L., Bernhardt, J., Huganir, R. L. (1996) Characterization of multiple phosphorylation sites on the AMPA receptor GluR1 subunit. *Neuron* **16**, 1179–1188
  55. Dumont, Y., Cadieux, A., Doods, H., Pheng, L. H., Abounader, R., Hamel, E., Jacques, D., Regoli, D., and Quirion, R. (2000) BIIIE0246, a potent and highly selective non-peptide neuropeptide YY(2) receptor antagonist. *Br. J. Pharmacol.* **129**, 1075–1088
  56. Toth, P. T., Bindokas, V. P., Bleakman, D., Colmers, W. F., and Miller, R. J. (1993) Mechanism of presynaptic inhibition by neuropeptide Y at sympathetic nerve terminals. *Nature* **364**, 635–639
  57. Klapstein, G. J., and Colmers, W. F. (1992) 4-Aminopyridine and low Ca<sup>2+</sup> differentiate presynaptic inhibition mediated by neuropeptide Y, baclofen and 2-chloroadenosine in rat hippocampal CA1 in vitro. *Br. J. Pharmacol.* **105**, 470–474
  58. Rhim, H., Kinney, G. A., Emmerson, P. J., and Miller, R. J. (1997) Regulation of neurotransmission in the arcuate nucleus of the rat by different neuropeptide Y receptors. *J. Neurosci.* **17**, 2980–2989
  59. Whittaker, E., Vereker, E., and Lynch, M. A. (1999) Neuropeptide Y inhibits glutamate release and long-term potentiation in rat dentate gyrus. *Brain Res.* **827**, 229–233
  60. Sanchez-Prieto, J., Budd, D. C., Herrero, I., Vazquez, E., Nicholls, D. G. (1996) Presynaptic receptors and the control of glutamate exocytosis. *Trends Neurosci.* **19**, 235–239
  61. Hori, T., Takai, Y., and Takahashi, T. (1999) Presynaptic mechanism for phorbol ester-induced synaptic potentiation. *J. Neurosci.* **19**, 7262–7267
  62. Tang, F. R., Lee, W. L., Gao, H., Chen, Y., Loh, Y. T., and Chia, S. C. (2004) Expression of different isoforms of protein kinase C in the rat hippocampus after pilocarpine-induced status epilepticus with special reference to CA1 area and the dentate gyrus. *Hippocampus* **14**, 87–98
  63. Hussain, R. J., and Carpenter, D. O. (2003) The effects of protein kinase C activity on synaptic transmission in two areas of rat hippocampus. *Brain Res.* **990**, 28–37
  64. Swartz, K. J., Merritt, A., Bean, B. P., Lovinger, D. M. (1993) Protein kinase C modulates glutamate receptor inhibition of Ca<sup>2+</sup> channels and synaptic transmission. *Nature* **361**, 165–168
  65. Lan, J. Y., Skeberdis, V. A., Jover, T., Grooms, S. Y., Lin, Y., Araneda, R. C., Zheng, X., Bennett, M. V., and Zukin, R. S. (2001) Protein kinase C modulates NMDA receptor trafficking and gating. *Nat. Neurosci.* **4**, 382–390
  66. Garcia-Sainz, J. A., Vazquez-Cuevas, F. G., and Romero-Avila, M. T. (2001) Phosphorylation and desensitization of alpha1-adrenergic receptors. *Biochem. J.* **353**, 603–610

Received for publication May 31, 2006.  
Accepted for publication September 29, 2006.